

Name:

DOB:

.....

Mild/moderate reaction:

- Swollen lips, face or eyes
- Itchy/tingling mouth
- Mild throat tightness
- Hives or itchy skin rash
- Abdominal pain or vomiting
- Sudden change in behaviour

Action to take:

- Stay with person, call for help if needed
- Locate adrenaline autoinjector(s)
- Give antihistamine:
.....

(If vomited, can repeat dose)

- Phone parent/emergency contact
- Do not take a shower to help with itchy skin, this can worsen the reaction

Emergency contact details:

1) Name:



2) Name:



Parental consent: I hereby authorise school staff to administer the medicines listed on this plan, in accordance with Department of Health Guidance on the use of AAI in schools.

Signed:

Print name:

Date:

Consent is required for children under 16 years (and for young people over 16 unable to give consent themselves) except in an unforeseen emergency

For more information about managing anaphylaxis in schools and "spare" adrenaline autoinjectors, visit: sparepensinschools.uk

Watch for signs of ANAPHYLAXIS

(a potentially life-threatening allergic reaction)

Anaphylaxis may occur without skin symptoms: ALWAYS consider anaphylaxis in someone with known food allergy who has **SUDDEN DIFFICULTY IN BREATHING**

A AIRWAY

- Persistent cough
- Hoarse voice
- Difficulty swallowing
- Swollen tongue

B BREATHING

- Difficult or noisy breathing
- Wheeze or persistent cough

C CONSCIOUSNESS

- Persistent dizziness
- Pale or floppy
- Suddenly sleepy
- Collapse/unconscious

IF ANY ONE (OR MORE) OF THESE SIGNS ABOVE ARE PRESENT:

1 Lie flat with legs raised (if breathing is difficult, allow person to sit)



2 Dial 999 for ambulance and say ANAPHYLAXIS ("ANA-FIL-AX-IS")

3 In a school with "spare" back-up adrenaline autoinjectors, **ADMINISTER** the **SPARE AUTOINJECTOR** if available

4 Stay with child/young person until ambulance arrives, **do NOT stand them up**

5 Phone parent/emergency contact. Phone parent/emergency contact. If you are on your own, call a friend or relative and ask them to come over.

6 Commence CPR if there are no signs of life

***** IF IN DOUBT, GIVE ADRENALINE *****

You can dial 999 from any phone, even if there is no credit left on a mobile. Medical observation in hospital is recommended after anaphylaxis. For more information about managing anaphylaxis in schools and "spare" back-up adrenaline autoinjectors, visit: sparepensinschools.uk

Additional instructions:

If wheezy due to an allergic reaction, DIAL 999 and GIVE ADRENALINE using a "back-up" adrenaline autoinjector if available, then use asthma reliever (e.g. blue puffer) via spacer, if prescribed

This BSACI Action Plan for Allergic Reactions is for children and young people with mild food allergies, who need to avoid certain allergens. For children/young adults at risk of anaphylaxis and who have been prescribed an adrenaline autoinjector device, there are BSACI Action Plans which include instructions for adrenaline autoinjectors. These can be downloaded at bsaci.org

For further information, consult NICE Clinical Guidance CG116 Food allergy in children and young people at guidance.nice.org.uk/CG116

This is a medical document to be completed by a healthcare professional. It must not be altered without their permission. This document provides medical authorisation for schools to administer a 'spare' back-up adrenaline autoinjector if needed, as permitted by the Human Medicines (Amendment) Regulations 2017. The healthcare professional named below confirms that there are no medical contra-indications to the above-named child being administered an adrenaline autoinjector by school staff in an emergency. **This plan has been prepared by:**

Sign & print name:

Hospital/Clinic:



Date: